

## New England Biolabs Certificate of Analysis

**Product Name:** Rapid PNGase F (non-reducing format)  
**Catalog #:** P0711S  
**Concentration:** 50 reactions  
**Unit Definition:** N/A  
**Lot #:** 0021709  
**Assay Date:** 09/2017  
**Expiration Date:** 9/2018  
**Storage Temp:** 4°C  
**Storage Conditions:** 50 mM NaCl, 20 mM Tris-HCl, 5 mM EDTA, (pH 7.5 @ 25°C)  
**Specification Version:** PS-P0711S v1.0  
**Effective Date:** 16 Feb 2016

Assay Name/Specification (minimum release criteria)	Lot #0021709
<p><b>Glycosidase Activity (Endo F1, F2, H)</b> - A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled Endo F1, F2, H substrate (Dansylated invertase high mannose) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (Endo F2, F3)</b> - A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled Endo F2, F3 substrate (Dansylated fibrinogen biantennary) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-Mannosidase)</b> - A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-Mannosidase substrate (Manβ1-4Manβ1-4Man-AMC) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (β-Xylosidase)</b> - A 10 µl reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled β-Xylosidase substrate (Xylβ1-4Xylβ1-4Xylβ1-4Xyl-AMC) and 1 µl of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>



## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0021709
<p><b>Glycosidase Activity (<math>\beta</math>1-3 Galactosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>1-4 Galactosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\beta</math>-Galactosidase substrate (Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>-N-Acetylgalactosaminidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\beta</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\beta</math>1-4Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\beta</math>-N-Acetylglucosaminidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\beta</math>-N-Acetylglucosaminidase substrate (GlcNAc<math>\beta</math>1-4GlcNAc<math>\beta</math>1-4GlcNAc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-Glucosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Glucosidase substrate (Glc<math>\alpha</math>1-6Glc<math>\alpha</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-Neuraminidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Neuraminidase substrate (Neu5Ac<math>\alpha</math>2-3Gal<math>\beta</math>1-3GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-2 Fucosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuc<math>\alpha</math>1-2Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Fucosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Fucosidase substrate (Fuc<math>\alpha</math>1-3Gal<math>\beta</math>1-4GlcNAc<math>\beta</math>1-3Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>




## New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0021709
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Galactosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-3Gal<math>\beta</math>1-4GlcNAc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-3 Mannosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-3Man<math>\beta</math>1-4GlcNAc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-6 Galactosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Galactosidase substrate (Gal<math>\alpha</math>1-6Gal<math>\alpha</math>1-6Glc<math>\alpha</math>1-2Fru-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>1-6 Mannosidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-Mannosidase substrate (Man<math>\alpha</math>1-6Man<math>\alpha</math>1-6(Man<math>\alpha</math>1-3)Man-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Glycosidase Activity (<math>\alpha</math>-N-Acetylgalactosaminidase)</b> - A 10 <math>\mu</math>l reaction in Rapid PNGase F Buffer (non-reducing format) containing 1 nM of fluorescently-labeled <math>\alpha</math>-N-Acetylgalactosaminidase substrate (GalNAc<math>\alpha</math>1-3(Fuc<math>\alpha</math>1-2)Gal<math>\beta</math>1-4Glc-AMC) and 1 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C results in no detectable activity as determined by thin layer chromatography.</p>	<b>Pass</b>
<p><b>Protease Activity (SDS-PAGE)</b> - A 20 <math>\mu</math>l reaction in 1X Rapid PNGase F Buffer (non-reducing format) containing 24 <math>\mu</math>g of a standard mixture of proteins and a minimum of 5 <math>\mu</math>l of Rapid PNGase F (non-reducing format) incubated for 20 hours at 37°C, results in no detectable degradation of the protein mixture as determined by SDS-PAGE with Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> - Rapid PNGase F (non-reducing format) is <math>\geq</math> 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>



Authorized by  
Derek Robinson  
16 Feb 2016



Inspected by  
Brad Landgraf  
25 Sep 2017

