

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: BamHI
Catalog Number: R0136M
Concentration: 100,000 U/mI

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Lot Number:10008861Expiration Date:05/2020Storage Temperature:-20°C

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0136T/M v1.0

BamHI Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0136MVIAL	BamHI	10008875	Pass	
B7203SVIAL	NEBuffer™ 3.1	0541804	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	0251805	Pass	

Assay Name/Specification	Lot # 10008861
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 60 Units of BamHl incubated for 4 hours at 37°C results in <10%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Blue-White Screening (Terminal Integrity)	Pass
A sample of pUC19 vector linearized with a 10-fold excess of BamHI, religated and	
transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	
Ligation and Recutting (Terminal Integrity)	Pass
After a 20-fold over-digestion of Lambda DNA with BamHI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,	
>95% can be recut with BamHI.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 20	
units of BamHI incubated for 16 hours at 37°C results in a DNA pattern free of	



R0136M / Lot: 10008861

Page 1 of 2

Assay Name/Specification	Lot # 10008861
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in NEBuffer 3.1 containing 1 µg of a mixture of single and	
double-stranded [3H] E. coli DNA and a minimum of 100 units of BamHI incubated for	
4 hours at 37°C releases <0.1% of the total radioactivity.	

This product has been tested and shown to be in compliance with all specifications.

Jianying Luo Production Scientist

18 May 2018

Michael Tonello

Packaging Quality Control Inspector

20 Jun 2018

