

New England Biolabs Certificate of Analysis

Product Name: AfIII
Catalog Number: R0520S
Concentration: 20,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of PhiX174 RF I DNA in rCutSmart™ Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.
Packaging Lot Number: 10205077
Expiration Date: 08/2025
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version: PS-R0520S/L v2.0


AfIII Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0520SVIAL	AfIII	10201746	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10198639	Pass
B6004SVIAL	rCutSmart™ Buffer	10198645	Pass

Assay Name/Specification	Lot # 10205077
Blue-White Screening (Terminal Integrity) A sample of Litmus28 vector linearized with a 10-fold excess of AfIII, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled pBR322 DNA and a minimum of 100 units of AfIII incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 100 units of AfIII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Functional Testing (15 minute Digest) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of PhiX174 RF I DNA and 1 µl	Pass

Assay Name/Specification	Lot # 10205077
of AfIII incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	
<p>Ligation and Recutting (Terminal Integrity) After a 5-fold over-digestion of PhiX174 DNA with AfIII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AfIII.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of PhiX174 RF I DNA and a minimum of 200 units of AfIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) AfIII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 20 units of AfIII is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



YunJie Sun
Production Scientist
10 Aug 2023



Josh Hersey
Packaging Quality Control Inspector
21 Aug 2023