

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Blpl
Catalog Number: R0585L
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10139760
Expiration Date: 11/2023
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 μg/ml BSA

Specification Version: PS-R0585S/L v1.0

Blpl Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0585LVIAL	Blpl	10126722	Pass	
B6004SVIAL	rCutSmart™ Buffer	10136926	Pass	

Assay Name/Specification	Lot # 10139760
Protein Purity Assay (SDS-PAGE)	Pass
Blpl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.	
Exonuclease Activity (Radioactivity Release)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and	
double-stranded [³H] E. coli DNA and a minimum of 100 units of Blpl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	
Thours at 37°C releases <0.1% or the total radioactivity.	
Endonuclease Activity (Nicking)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and	
a minimum of 30 units of Blpl incubated for 4 hours at 37°C results in <20%	
conversion to the nicked form as determined by agarose gel electrophoresis.	
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of	
10 units of Blpl incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	
Ligation and Recutting (Terminal Integrity)	Pass



R0585L / Lot: 10139760

Page 1 of 2



Assay Name/Specification	Lot # 10139760
After a 10-fold over-digestion of Lambda DNA with BlpI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with BlpI.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

01 Mar 2022

Michael Tonello

Packaging Quality Control Inspector

01 Mar 2022

