

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Blpl  |
|------------------------|---|
| Catalog Number:        | R0585S  |
| Concentration:         | 10,000 U/ml   |
| Unit Definition:       | One unit is defined as the amount of enzyme required to digest 1 μg<br>of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl. |
| Packaging Lot Number:  | 10160142  |
| Expiration Date:       | 06/2024   |
| Storage Temperature:   | -20°C   |
| Storage Conditions:    | 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%<br>Glycerol, 200 μg/ml BSA  |
| Specification Version: | PS-R0585S/L v1.0  |

| Blpl Component List    |                       |            |                      |  |
|------------------------|-----------------------|------------|----------------------|--|
| <b>NEB Part Number</b> | Component Description | Lot Number | Individual QC Result |  |
| R0585SVIAL             | Blpl                  | 10154077   | Pass                 |  |
| B6004SVIAL             | rCutSmart™ Buffer     | 10158558   | Pass                 |  |

| Assay Name/Specification  | Lot # 10160142 |
|---|----------------|
| Endonuclease Activity (Nicking)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of supercoiled PhiX174 DNA and<br>a minimum of 30 units of Blpl incubated for 4 hours at 37°C results in <20%<br>conversion to the nicked form as determined by agarose gel electrophoresis.            | Pass           |
| Exonuclease Activity (Radioactivity Release)<br>A 50 µl reaction in CutSmart <sup>™</sup> Buffer containing 1 µg of a mixture of single and<br>double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of BlpI incubated for 4<br>hours at 37°C releases <0.1% of the total radioactivity. | Pass           |
| <b>Ligation and Recutting (Terminal Integrity)</b><br>After a 10-fold over-digestion of Lambda DNA with Blpl, >95% of the DNA fragments<br>can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments,<br>~75% can be recut with Blpl.   | Pass           |
| <b>Protein Purity Assay (SDS-PAGE)</b><br>Blpl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.  | Pass           |
| Non-Specific DNase Activity (16 Hour)   | Pass           |





be INSPIRED drive DISCOVERY stay GENUINE

240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

| Assay Name/Specification   | Lot # 10160142 |
|--|----------------|
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 10 units of BlpI incubated for 16 hours at 37⁰C results in a DNA pattern free of |                |
| detectable nuclease degradation as determined by agarose gel electrophoresis.  |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang

Production Scientist 08 Sep 2022

Erin Varney

Packaging Quality Control Inspector 08 Sep 2022

