

ERY 240 County Road E Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name:BlplCatalog Number:R0585SConcentration:10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10172474
Expiration Date: 12/2024
Storage Temperature: -20°C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 200 µg/ml BSA

Specification Version: PS-R0585S/L v1.0

| Blpl Component List | | | | |
|------------------------|-----------------------|------------|----------------------|--|
| NEB Part Number | Component Description | Lot Number | Individual QC Result | |
| R0585SVIAL | Blpl | 10172473 | Pass | |
| B6004SVIAL | rCutSmart™ Buffer | 10173160 | Pass | |

| Assay Name/Specification | Lot # 10172474 |
|--|----------------|
| Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of | Pass |
| 10 units of Blpl incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | |
| Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart™ Buffer containing 1 μg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of Blpl incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Protein Purity Assay (SDS-PAGE) Blpl is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection. | Pass |
| Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with BlpI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, ~75% can be recut with BlpI. | Pass |
| Endonuclease Activity (Nicking) | Pass |



R0585S / Lot: 10172474

Page 1 of 2



| Assay Name/Specification | Lot # 10172474 |
|---|----------------|
| A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 30 units of Blpl incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis. | |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Suń Production Scientist

10 Jan 2023

Michael Tonello

Packaging Quality Control Inspector 13 Jan 2023



R0585S / Lot: 10172474

Page 2 of 2