

## New England Biolabs Certificate of Analysis

**Product Name:** Sspl-HF®  
**Catalog Number:** R3132M  
**Concentration:** 100,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10159244  
**Expiration Date:** 08/2024  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)  
**Specification Version:** PS-R3132M v2.0

Sspl-HF® Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3132MVIAL	Sspl-HF®	10159242	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10156427	Pass
B6004SVIAL	rCutSmart™ Buffer	10156426	Pass

Assay Name/Specification	Lot # 10159244
<b>Protein Purity Assay (SDS-PAGE)</b> Sspl-HF® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 20 units of Sspl-HF® is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 100 units of Sspl-HF® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Non-Specific DNase Activity (16 Hour)</b>	Pass

Assay Name/Specification	Lot # 10159244
<p>A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of SspI-HF® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	
<p><b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of Lambda DNA with SspI-HF®, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with SspI-HF®.</p>	<b>Pass</b>
<p><b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of SspI-HF® incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



\_\_\_\_\_  
Penghua Zhang  
Production Scientist  
19 Aug 2022



\_\_\_\_\_  
Michael Tonello  
Packaging Quality Control Inspector  
19 Aug 2022