

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name:	Agel-HF®
Catalog Number:	R3552S
Concentration:	20,000 U/ml
Unit Definition:	One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction of 50 μl.
Packaging Lot Number:	10235830
Expiration Date:	02/2026
Storage Temperature:	-20°C
Storage Conditions:	10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version:	PS-R3552S/L v2.0

Agel-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3552SVIAL	Agel-HF®	10225346	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10234872	Pass	
B6004SVIAL	rCutSmart™ Buffer	10235241	Pass	

Assay Name/Specification	Lot # 10235830
<b>Blue-White Screening (Terminal Integrity)</b> A sample of LITMUS28i vector linearized with a 10-fold excess of AgeI-HF®, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 200 units of AgeI-HF® incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in rCutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and 1 µl of Agel-HF® incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with AgeI-HF®, >95% of the DNA	Pass





240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Assay Name/Specification	Lot # 10235830
fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with AgeI-HF®.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in rCutSmart <sup>™</sup> Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of Agel-HF® incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) AgeI-HF® is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 20 units of AgeI-HF® is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

YunJie Sun Production Scientist 23 Jan 2024

Michael 11.

Michael Tonello Packaging Quality Control Inspector 22 Apr 2024

