

## New England Biolabs Certificate of Analysis

**Product Name:** BsrGI-HF  
**Catalog #:** R3575S/L  
**Concentration:** 20,000 units/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Lot #:** 0021506  
**Assay Date:** 06/2015  
**Expiration Date:** 06/2017  
**Storage Temp:** -20°C  
**Storage Conditions:** 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 µg/ml BSA  
**Specification Version:** PS-R3575S/L v1.0  
**Effective Date:** 27 Oct 2014

Assay Name/Specification (minimum release criteria)	Lot #0021506
<b>Endonuclease Activity (Nicking)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of supercoiled ΦX174 DNA and a minimum of 60 units of BsrGI-HF incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Exonuclease Activity (Radioactivity Release)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] <i>E. coli</i> DNA and a minimum of 100 units of BsrGI-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	<b>Pass</b>
<b>Functional Test (15 minute Digest)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of BsrGI-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Ligation and Recutting (Terminal Integrity)</b> - After a 10-fold over-digestion of Lambda DNA with BsrGI-HF, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsrGI-HF.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 Hour)</b> - A 50 µl reaction in CutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of BsrGI-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Protein Purity Assay (SDS-PAGE)</b> - BsrGI-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	<b>Pass</b>



---


## New England Biolabs Certificate of Analysis

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



---

Authorized by  
Derek Robinson  
27 Oct 2014



---

Inspected by  
Toby Claus  
18 Jun 2015

