

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Streptavidin Magnetic Beads

Catalog Number: \$1420\$
Concentration: 4 mg/ml
Packaging Lot Number: 10080123
Expiration Date: 09/2022
Storage Temperature: 4°C

Storage Conditions: 0.05 % NaN3, 0.1 % BSA, 0.05 % Tween®20, 1 X PBS, (pH 7.4 @ 25°C)

Specification Version: PS-S1420S v1.0

Streptavidin Magnetic Beads Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
S1420SVIAL	Streptavidin Magnetic Beads	10051761	Pass	

Assay Name/Specification	Lot # 10080123
Binding Capacity (Magnetic Beads) Streptavidin Magnetic Beads ( 500 μg ) were equilibrated and incubated with 100 μl of 5 μM 5'-Biotin-dT25-FAM-3' for 1 hour at 25°C. Binding capacity was determined to	Pass
be >500 pmol of oligo per mg of beads.	
Functional Binding Assay (Qualitative) Streptavidin Magnetic Beads ( 500 μg ) were equilibrated and incubated with 200 μl of Biotin Mouse Anti-Human IgG then washed and incubated with 500 μl Human Serum IgG for 1 hour at 25°C, then washed, eluted and evaluated by Tris-Glycine gel to confirm low non-specific binding of extract proteins and high isolation of target.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in Streptavidin Magnetic Bead Storage Buffer containing 1 µg of PhiX174-HaeIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in Streptavidin Magnetic Bead Storage Buffer containing 1 µg of PhiX174-HaelII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
RNase Activity (Buffer) A 10 µl reaction in Streptavidin Magnetic Bead Storage Buffer containing 40 ng of a	Pass



S1420S / Lot: 10080123

Assay Name/Specification	Lot # 10080123
300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours,	
>90% of the substrate RNA remains intact as determined by fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Michael Sproviero-

Production Scientist

12 Aug 2020

Josh Hersey

Packaging Quality Control Inspector

12 Aug 2020



S1420S / Lot: 10080123

Page 2 of 2