

New England Biolabs Product Specification

<i>Product Name:</i>	<i>Bst DNA Polymerase, Large Fragment</i>
<i>Catalog #:</i>	<i>M0275M</i>
<i>Concentration:</i>	<i>120,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 65°C.</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.1 % Triton®X-100 , 50 % Glycerol, (pH 7.1 @ 25° C)</i>
<i>Specification Version:</i>	<i>PS-M0275M v1.0</i>
<i>Effective Date:</i>	<i>04 Aug 2015</i>

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment incubated for 4 hours at either 37°C or 65°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment incubated for 4 hours at either 37°C or 65°C releases <0.1% of the total radioactivity.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 120 units of *Bst* DNA Polymerase, Large Fragment incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of Lambda DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 500 units of *Bst* DNA Polymerase, Large Fragment incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Phosphatase Activity (pNPP) - A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM *p*-Nitrophenol Phosphate (pNPP) and a minimum of 100 units *Bst* DNA Polymerase, Large Fragment incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

Protein Purity Assay (SDS-PAGE) - *Bst* DNA Polymerase, Large Fragment is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)

qPCR DNA Contamination (*E. coli* Genomic) - A minimum of 120 units of *Bst* DNA Polymerase, Large Fragment is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.

RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of *Bst* DNA Polymerase, Large Fragment is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.



Derek Robinson
Director of Quality Control

Date 04 Aug 2015

