

New England Biolabs Product Specification

<i>Product Name:</i>	<i>OneTaq[®] Hot Start Quick-Load[®] 2X Master Mix with GC Buffer</i>
<i>Catalog #:</i>	<i>M0489S/L</i>
<i>Concentration:</i>	<i>2X Concentrate</i>
<i>Shelf Life:</i>	<i>24 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Composition (1X):</i>	<i>80 mM Tris-SO₄ (pH 9.2 @ 25°C), 20 mM (NH₄)₂SO₄, 2 mM MgSO₄, 0.2 mM dATP, 0.2 mM dCTP, 0.2 mM dGTP, 0.2 mM dTTP, 5 % Glycerol, 5 % DMSO, 0.06 % IGEPAL[®] CA-630, 0.05 % Tween[®] 20, 1 X Xylene cyanol, 1 X Tartrazine, 25 units/ml OneTaq[®] Hot Start DNA Polymerase</i>
<i>Specification Version:</i>	<i>PS-M0489S/L v2.0</i>
<i>Effective Date:</i>	<i>12 Feb 2020</i>

Assay Name/Specification (minimum release criteria)

Inhibition of Primer Extension (Hot Start, Radioactivity Incorporation) - A 50 µl primer extension assay in ThermoPol[®] Reaction Buffer in the presence of 200 µM dNTPs including [³H]-dTTP, containing 15 nM primed single-stranded M13mp18 with 2.5 units of OneTaq[®] Hot Start DNA Polymerase incubated for 16 hours at 25°C yields >95% inhibition when compared to a non-hot start control reaction.

Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in 1X OneTaq[®] Hot Start Quick-Load[®] Master Mix with GC Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

PCR Amplification (Buffer Dependent, >65% GC-rich, Master Mix) - A 25 µl reaction in 1X OneTaq[®] Hot Start Quick-Load[®] Master Mix with GC Buffer and 0.2 µM primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the buffer-dependent production of the 737 bp product.

PCR Amplification (Enhancer Dependent, >70% GC-rich, Master Mix) - A 25 µl reaction in 1X OneTaq[®] Hot Start Quick-Load[®] Master Mix with GC Buffer and 20% OneTaq[®] High GC Enhancer in the presence of 0.2 µM primers containing 10 ng Human Genomic DNA for 30 cycles of PCR amplification results in the enhancer-dependent production of the 627 bp product.

PCR Amplification (Hot Start 2 kb Lambda DNA) - A 25 µl reaction in OneTaq[®] Standard Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 10 pg Lambda DNA and 50 ng Human Genomic DNA with 0.625 units of OneTaq[®] Hot Start DNA Polymerase for 30 cycles of PCR amplification results in an increase in yield of the 2 kb Lambda product and a decrease in non-specific genomic bands when compared to a non-hot start control reaction.



New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)

RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of One Taq [®] Hot Start Quick-Load [®] 2X Master Mix with GC Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 12 Feb 2020

Derek Robinson
Director, Quality Control

